

Recent Progresses of the DAMPE Mission*

CHANG Jin

(Key Laboratory of Dark Matter and Space Astronomy, Purple Mountain Observatory, Chinese Academy of Sciences, Nanjing 210033)

(School of Astronomy and Space Science, University of Science and Technology of China, Hefei 230026)

Abstract The DArk Matter Particle Explorer (DAMPE) is a space high-energy particle and γ -ray detector whose major scientific goals are the indirect detection of dark matter particles, the origin of cosmic rays and high-energy γ -ray astronomy. Since its successful launch in December 2015, the DAMPE has been operated smoothly in orbit for more than 8 years. The direct measurements of the boron-to-carbon and boron-to-oxygen flux ratios, and the proton+ Helium spectrum up to 316 TeV have been obtained, revealing new spectral features with very high significances. The search results of γ -ray spectral line and fractionally charged particles indicate a good potential of DAMPE for indirect dark matter detection and new physical discovery in space. The DAMPE measurements are expected to significantly advance our understanding of the fundamental problems in astroparticle physics.

Key words DAMPE, Dark matter, Indirect detection, Cosmic rays

Classified index P35, P1

1 Introduction

It has been well established that our Universe is made up of about 4.9% baryonic matter, 26.8% Dark Matter (DM), and 68.3% dark energy^[1]. The physical nature of DM is one of the most important fundamental questions of modern physics. Many types of candidate DM models have been proposed, among which the leading one is a kind of Weakly Interacting Massive Particles (WIMPs) beyond the standard model of particle physics, as motivated by the relic abundance of DM and the bottom-up evolution pattern of the large scale structures of the Universe^[2,3]. Many efforts have been made to search for the WIMP DM in laboratories, including the direct detection of the WIMP-nucleon scattering by underground detectors, the indirect detection of the annihilation or decay products in the cosmic radiation, and the collider detection of WIMP pairs produced in high-energy parti-

cle colliders^[4]. After several decades of efforts, no convincing evidence has been found yet. Particularly, the direct detection experiments push the WIMP-nucleon scattering strength to a very low level^[5], which has already severely constrained a series of WIMP models.

In recent years, some interesting anomalies have been found in Cosmic Ray (CR) and γ -ray observations, such as the positron excess discovered by PAMELA^[6], Fermi-LAT^[7], and AMS-02^[8], and the associated total Cosmic Ray Electron plus positron (CRE) excess by AT-IC^[9], Fermi-LAT^[10], and AMS-02^[11]. The excesses of positrons and CREs can be explained by either the astrophysical sources such as pulsars and the DM annihilation or decay^[12–17]. There are still debates on the astrophysical or DM origin of the positron and/or CRE excesses, in particular, that the most recent HAWC observations of two nearby pulsars tend to disfavor the pulsar interpretation^[18]. Higher precision measurements

* Supported by the National Key Research and Development Program of China (2016 YFA0400200), and the National Natural Science Foundation of China (U1738205, U1738206)

Received May 15, 2024

©The Author(s) 2024. This is an open access article under the CC-BY 4.0 License (<https://creativecommons.org/licenses/by/4.0/>)

in a wider energy band are crucial to identifying their physical origin. Other anomalies include the γ -ray excess in the Galactic center^[19], and the potential excess of CR antiprotons^[20,21].

The purpose of the DArk Matter Particle Explorer (DAMPE) is to search for possible annihilation/decay relics of DM in CREs, with unprecedentedly high energy resolution and low background^[22,23]. DAMPE is also proposed as a high-energy particle detector for observations of γ -rays and CR nuclei, which can be particularly helpful in the study of high-energy astrophysical phenomena and CR physics^[23]. The DAMPE mission was launched into a 500 km Sun-synchronous orbit on 17 December 2015, with an inclination angle of 97°. It orbits the Earth every 95 minutes. All the detectors perform excellently in space. The daily event rate of DAMPE is about 5 million. It has recorded in total over 8 billion CR events as of May 2020, most of which are CR nuclei. There are about 1% CREs and 3×10^{-5} γ -ray photons in the DAMPE data.

2 Detector

DAMPE is made of 4 sub-detectors, which are, from top to bottom, the Plastic Scintillator strip Detector (PSD^[24]), the Silicon-Tungsten trackKer-converter (STK^[25]), the BGO imaging calorimeter^[26], and the NeUtron Detector (NUD^[27]). A schematic plot of the DAMPE detector is shown in Figure 1^[23]. The total weight of the payload (satellite) is about 1.5 (1.9) tons, and the power consumption is 300 (500) W. The size of the satellite is about 1.2 m $\times$ 1.2 m $\times$ 1.0 m.

The PSD is to measure the (absolute) charge of incident particles up to $z=30$ via the ionization process

(dE/dx). It also serves as an anticoincidence detector for γ -rays. The PSD consists of two layers of orthogonally placed plastic scintillator bars. For each layer, 41 plastic scintillator bars are further placed in two sub-layers with a shift of 0.8 cm between them to reduce the gaps. The effective area of the PSD is 82.5 cm $\times$ 82.5 cm^[23]. The weight of the PSD is about 103 kg, and the power is 8.5 W.

The main function of STK is to measure the trajectory of a particle. The charge of light nuclei ($z < 8$) can also be measured by the STK. The STK consists of 12 layers (6 for x direction and 6 for y direction) of silicon trackers, each with two sub-layers arranged orthogonally to measure the x and y positions of a track. The STK is also a γ -ray converter by means of three 1 mm thick tungsten plates inserted in the second, third, and fourth planes. Each silicon layer is made of 16 ladders, each formed by 4 silicon micro-strip detectors. Every ladder is segmented into 768 strips, half of which are readout strips and the other half are floating strips. In total, there are $384 \times 16 \times 12 = 73728$ readout channels of the STK. The effective detection area of STK is 76 cm $\times$ 76 cm. The STK has a weight of 154 kg and a power consumption of 82 W.

The BGO calorimeter is the major instrument of DAMPE. It has three functions: (i) measuring the energy deposition of incident particles, (ii) imaging the 3D profile of the shower development, and providing electron/hadron discrimination, and (iii) providing the level 0 trigger for the DAMPE data acquisition system. The BGO calorimeter is made of 308 BGO bars, each is read out at two ends by PhotoMultiplier Tubes (PMTs). The BGO crystals are placed orthogonally in 14 layers, with a total depth of about 32 radiation lengths and about




Fig. 1 Schematic plot of the DAMPE detectors

1.6 nuclear interaction lengths. The length of the BGO crystal is 60 cm, which is the longest in the world. The calorimeter has an effective area of $60 \text{ cm} \times 60 \text{ cm}$, a total weight of 1052 kg, and a power consumption of 42 W.

The NUD is at the bottom of the detector. The main purpose of the NUD is to perform electron/hadron discrimination by means of the fact that neutrons are much more abundant in hadronic showers than in electromagnetic ones. Once the neutrons are created, they are quickly thermalized in the BGO calorimeter, and then get captured by Boron-doped plastic scintillators. The NUD is made of four $30 \text{ cm} \times 30 \text{ cm} \times 1 \text{ cm}$ plastic scintillators with 5% (weight) Boron element. The NUD has an effective area of $61 \text{ cm} \times 61 \text{ cm}$, a mass of 12 kg, and a power of 0.5 W.

Table 1 summarizes the expected detector performance of DAMPE, for observations of electrons, γ -rays, and protons^[23]. With good measurements of the charge, direction, and energy of various types of particles, DAMPE is expected to play a significant role in exploring the high-energy Universe.

3 On-orbit Performance

Dedicated calibration of each unit of the detector has been done using the on-orbit data, which enables precise measurements of the physical quantities of each event^[28]. Figure 2 gives the charge measurements by the PSD^[29]. The absolute charge resolution is about 0.20e (e means unit charge) for $z=8$ (Oxygen) and 0.30e for $z=26$ (Iron). Such a good charge resolution is the basis of precise spectral measurements of each species. Using the

bright γ -ray sources we can calibrate the angular resolution of the STK, which results in about 0.5° for energies higher than 5 GeV.

The very thick BGO calorimeter enables no significant leakage of electromagnetic showers from the bottom for energies as high as TeV. There is, however, a few percents of energy loss due to the dead materials of the calorimeter. An event-level energy correction method was developed and verified by MC simulations and the beam test data^[30]. The beam test data and the MC simulations show that the energy resolution for CREs is better than 1.2% for energies above 100 GeV. DAMPE also has P- and N-side readouts for each BGO crystal, which have different gains. These two-side readouts give a consistency check of the energy measurement. We find that the energies measured by both sides agree with each other very well. Figure 3 presents the ratios of energies measured by the two sides, together




Fig. 2 Charge spectrum obtained using two years of on-orbit data




Fig. 3 Ratios of energies reconstructed with P- and N-side data

Table 1 Expected performance of DAMPE

Parameter	Value
Energy range (e/ γ)	5 GeV to 10 TeV
Energy resolution (e/ γ)	1.5% at 800 GeV
Energy range (p)	50 GeV to 500 TeV
Energy resolution (p)	40% at 800 GeV
Effective area (vertical γ)	1100 cm^2 at 100 GeV
Geometry factor (e)	$0.3 \text{ m}^2 \cdot \text{sr}$ above 30 GeV
Angular resolution (γ)	0.1° at 100 GeV
Field of view	1.0 sr

with a Gaussian fit that gives a mean of 1.005 ± 0.005 and a width of 0.016 ± 0.001 ^[31]. Such a result further supports the quoted energy resolution of about 1%.

Particle identification is one of the most challenging things for the observations of CREs. The background of CREs is CR protons (heavier nuclei can be effectively rejected *via* the charge measurement in PSD). The fluxes of CREs are about three orders of magnitude lower than that of protons. Therefore, a high rejection power of protons is essential for reliable measurements of the CRE fluxes. DAMPE uses the morphology difference between CREs and protons in the calorimeter to distinguish them^[32,33]. This is basically due to different developments of hadronic and electromagnetic showers in the calorimeter. A simple two-parameter method was adopted to describe the longitude and latitude developments of the showers. They are: (i) an energy-weighted Root-Mean-Square (RMS) value of hit positions in the calorimeter (shower spread, L_{ss}), and (ii) the deposited energy fraction of the last BGO layer (F_{last}). Figure 4 (a) shows the distributions of these two parameters for the flight data with BGO energies between 500 GeV and 1 TeV^[31]. We can see clearly two populations of events in this plot, with the upper population being proton candidates and the lower one being CRE candidates. To better quantify the separation between these two populations, we further define a shape parameter, $\zeta = F_{last} \times (L_{ss}/\text{mm})^4 / (8 \times 10^6)$, which projects these two parameters into one dimension. The distributions of the ζ parameter for the flight data and Monte Carlo (MC) simulations for CREs and protons are shown in the right panel of Figure

5. We select CREs with $\zeta \leq 8.5$, which gives only about 2% proton background in this energy range^[31]. For energies below TeV, the proton background is lower than 3%. The highest background in the energy range from 25 GeV to 4.6 TeV is estimated to be about 18%^[31].

The NUD provides additional help of the electron-proton discrimination, since the neutron yields for hadronic showers are much higher than those of the electromagnetic showers. Simulations show that using the NUD alone the proton background can be suppressed by a factor of about 25 when keeping a CRE efficiency of 95%, for 1–5 TeV deposited energies^[34]. Together with the calorimeter, the total suppression power improves by a factor of about 2^[34]. The NUD capability is expected to be more important at higher energies.

4 Scientific Results

Figure 4 shows the direct measurements of Boron-to-Carbon (B/C) and Boron-to-Oxygen (B/O) flux ratios in the energy range from 10 GeV/n (GeV per nucleon) to 5.6 TeV/n with 6 years of on-orbit data collected by DAMPE, with high statistics and well-controlled systematic uncertainties^[35]. Boron nuclei in cosmic rays are believed to be mainly produced by the fragmentation of heavier nuclei, such as carbon and oxygen, *via* collisions with the interstellar matter. Therefore, the precise measurements of the B/C and B/O ratios are very essential for probes of the CR propagation. The energy dependence of both the B/C and B/O ratios measured by DAMPE can be well fitted by a broken power-law model rather than a single power-law model, suggesting




Fig. 4 Boron-to-carbon (a) and boron-to-oxygen (b) flux ratios as functions of kinetic energy per nucleon measured by DAMPE^[35] (red filled dots), compared with measurements from other experiments^[37–44]

the existence in both flux ratios of a spectral hardening at about 100 GeV/n. The significance of the break is about 5.6σ and 6.9σ for the GEANT4 simulation, and 4.46σ and 6.9σ for the alternative FLUKA simulation, for B/C and B/O, respectively. These latest measurements deviate from the predictions of conventional turbulence theories of the Interstellar Medium (ISM), which point toward a change of turbulence properties of the ISM at different scales or novel propagation effects of CRs, and should be properly incorporated in the indirect detection of dark matter *via* anti-matter particles. Improved measurements of the B/C, B/O, and other secondary-to-primary ratios in cosmic rays with higher statistics and lower systematics by DAMPE and future direct detection experiments such as HERD^[36] are expected to eventually uncover the fundamental problems of the origin

and propagation of CRs and shed new light on the indirect detection of dark matter particles.

Figure 6 shows the direct measurement of the CR proton+Helium (p+He) spectrum from 46 GeV to 316 TeV with 6 years of on-orbit data collected by DAMPE^[45]. The selection of proton+helium, instead of individual proton and helium contributions, allows the collection of additional statistics, thus reaching higher energies with low background. The DAMPE p+He spectrum is in agreement with other direct/indirect-detection experiments, within the systematic uncertainties. The DAMPE measurement thus provides a link between direct and indirect CR measurements, exhibiting a good general agreement among very different techniques. The p+He spectrum in such a wide energy range deviates from a simple power-law behavior, and shows the pres-




Fig. 5 (a) Shower spread versus the last layer energy fraction for selected events with BGO energies between 500 GeV and 1 TeV. (b) One-dimensional distributions of the shape parameter ζ , compared with MC simulations




Fig. 6 P+He spectrum from 46 GeV to 316 TeV measured by DAMPE^[45] (red fill dots), compared with direct measurements of p+He made by ATIC-02^[48], NUCLEON^[49] and CREAM^[50] (a), and indirect measurements from ARGO-YBJ+WFCT^[51], HAWC^[52], KASCADE^[53] and EAS-TOP+MACRO^[54] (b)

ence of a spectral hardening around about 600 GeV followed by a softening at 28.8 ± 4.5 TeV. The softening features revealed in both the individual proton^[46] and helium^[47] spectra measured by DAMPE are further confirmed by the p+He spectrum, with an unprecedented significance of 6.6σ . Moreover, The p+He spectrum shows a novel hint regarding a possible direct detection of a second hardening around 150 TeV, for the first time in space.

Figure 7 shows the results of the Fractionally Charged Particle (FCP) search based on the on-orbit data of the DAMPE experiment^[55]. Unlike underground experiments, which require an FCP energy of the order of hundreds of GeV, the FCP search with DAMPE starts at only a few GeV. Based on the analysis using 5 years of cosmic-ray events recorded by DAMPE, no FCP signal is observed (Figure 7a) and the flux upper limit of $\frac{2}{3}e$ FCP is found to be $\Phi < 6.2 \times 10^{-10} \text{ cm}^{-2} \cdot \text{sr}^{-1} \cdot \text{s}^{-1}$ at the 90% CL (Confidence Level, see Figure 7b). Our results demonstrate that DAMPE exhibits higher sensitivity than experiments of similar types by three orders of magnitude that more stringently restricts the conditions for the existence of FCP in primary cosmic rays. In the future, with the accumulation of more on-orbit data, DAMPE is expected to perform even more sensitive FCP searches.

Thanks to the unprecedented high energy resolution, DAMPE is well-suitable for searching for monochromatic and sharp γ -ray structures in the energy range from GeV to TeV. With the first 5 years of

DAMPE data, we carry out the search for spectral γ -ray lines from 10 to 300 GeV^[63]. To improve the sensitivity, we develop two types of dedicated data and adopt the signal-to-noise ratio optimized Regions of Interest (ROIs) for different DM density profiles. No line signals or candidates with TS value ≥ 9 are detected between 10 and 300 GeV in the Galaxy. The 95% CL constraints on the annihilation cross section for $\chi\chi \rightarrow \gamma\gamma$ and the decay lifetime for $\chi \rightarrow \gamma\nu$, with proper addressing of the systematic uncertainties, are presented in Figure 8. Most of our constraints are comparable to the 5.8-year results of Fermi-LAT thanks to the better energy resolution and the smaller influence of the systematic uncertainties. For the decaying DM, the DAMPE lower limits on the decay lifetime are stronger for DM with mass ≤ 100 GeV by a factor of about 2. In view of the fact that the DAMPE data set is about ten times smaller than that of Fermi-LAT, our findings demonstrate the potential of high-energy-resolution observations on dark matter detection.

Figure 9 shows the all-sky γ -ray map above 2 GeV along with the identified point-like sources obtained using 7.2 years of the DAMPE data^[65]. The luminous Galactic plane and some bright sources can be clearly seen. Using the blind search method based on the Li-Ma formula^[66], 248 point sources in total are detected with TS values larger than 25 and their locations are shown in Figure 10. By associating the observed sources with the Fermi-LAT 4 FGL-DR3 catalog^[67], the types of these sources are determined. The analysis reveals that a significant portion of the sources are associated with Ac-




Fig. 7 (a) Distribution of PSD-STK charge for the on-orbit data of DAMPE, the red lines indicate the signal region for FCPs. (b) FCP flux upper limit of DAMPE^[55] (red dots), compared with the other space experiments^[56,57] and underground experiments^[58–62]




Fig. 8 95% CL constraints for different DM density profiles with the DAMPE experiment^[63]. Blue dot-dashed lines show the 5.8-year Fermi-LAT constraints^[64]




Fig. 9 Significant point sources ($TS \geq 25$) detected in 7.2-year DAMPE data^[65]. Markers present the types of the associated sources. The numbers of different source types are listed in the right table

tive Galactic Nuclei (AGNs) and pulsars, indicating their origin in high-energy astrophysical processes associated with these objects.

Figure 10 shows the preliminary spectral measurement of the Galactic Center Excess (GCE) by DAMPE and the preferred DM parameter space for the annihila-

tion channel $\chi\chi \rightarrow b\bar{b}$ ^[68]. The GCE is searched with 7.2 years of the DAMPE photon data. It is detected with a TS value of 81.3, which corresponds to a significance of 7.9σ with 6 degrees of freedom. The spatial distribution is fitted with the DM annihilation distribution of the generalized NFW profile $\rho_{\text{NFW}}(r) = \rho_0 / [(r/r_s)^\gamma]$




Fig. 10 Preliminary spectral measurement of the Galactic center excess (GCE) by DAMPE (a) and the preferred DM parameter space for the annihilation channel $\chi\chi \rightarrow b\bar{b}$ (b)^[68]

$(1 + r/r_s)^{3-\gamma}]$ and the optimal inner slope is $\gamma = 1.16 \pm 0.10$, which is consistent with that of the Fermi-LAT^[69]. If the excess is explained with the DM annihilation into $b\bar{b}$, DM particles with mass $m_\chi = 45 \pm 11$ GeV and cross section $\langle \sigma v \rangle = (2.2 \pm 0.3) \times 10^{-26} \text{ cm}^3 \cdot \text{s}^{-1}$ is required. The DM parameter space is also consistent with the Fermi-LAT^[70,71].

5 Summary

DAMPE has been operating smoothly in orbit for more than eight years. Based on the accumulated high-quality data recorded by DAMPE, multiple latest scientific results have been obtained, regarding CR physics, indirect DM detection and γ -ray astronomy. The precise measurements of the B/C and B/O ratios reveal a common spectral hardening at about 100 GeV/n with very high significances, which provides important implications in understanding the CR propagation. The latest measurements of the p+He spectrum up to 316 TeV suggest a hint of hardening structure at about 150 TeV for the first time in space. The FCP search provides the strictest restriction in space for the existence of FCP in primary cosmic rays. Benefiting from the unprecedented high energy resolution of DAMPE, the γ -ray line search achieves even stronger DM constraints for the lower limits on the decay lifetime below 100 GeV. In the near future, more scientific progresses from DAMPE are expected to shed new light on our understandings of DM and the high-energy Universe.

References

- [1] ADE P A R, AGHANIM N, ARNAUD M, *et al.* Planck 2015 results. XIII. Cosmological parameters[J]. *Astronomy & Astrophysics*, 2016, **594**: A13
- [2] JUNGMAN G, KAMIONKOWSKI M, GRIEST K. Supersymmetric dark matter[J]. *Physics Reports*, 1996, **267**(5/6): 195-373
- [3] BERTONE G, HOOPER D, SILK J. Particle dark matter: evidence, candidates and constraints[J]. *Physics Reports*, 2005, **405**(5/6): 279-390
- [4] BI X J, YIN P F, YUAN Q. Status of dark matter detection[J]. *Frontiers of Physics*, 2013, **8**(6): 794-827
- [5] ZHAO L, LIU J L. Experimental search for dark matter in China[J]. *Frontiers of Physics*, 2020, **15**(4): 44301
- [6] ADRIANI O, BARBARINO G C, BAZILEVSKAYA G A, *et al.* An anomalous positron abundance in cosmic rays with energies 1.5-100 GeV[J]. *Nature*, 2009, **458**(7238): 607-609
- [7] ACKERMANN M, AJELLO M, ALLAFORT A, *et al.* Measurement of separate cosmic-ray electron and positron spectra with the Fermi Large Area Telescope[J]. *Physical Review Letters*, 2012, **108**(1): 011103
- [8] AGUILAR M, ALBERTI G, ALPAT B, *et al.* First result from the Alpha Magnetic Spectrometer on the International Space Station: precision measurement of the positron fraction in primary cosmic rays of 0.5-350 GeV[J]. *Physical Review Letters*, 2013, **110**(14): 141102
- [9] CHANG J, ADAMS J H, AHN H S, *et al.* An excess of cosmic ray electrons at energies of 300-800 GeV[J]. *Nature*, 2008, **456**(7220): 362-365
- [10] ABDO A A, ACKERMANN M, AJELLO M, *et al.* Measurement of the cosmic ray $e^+ + e^-$ spectrum from 20 GeV to 1 TeV with the Fermi Large Area telescope[J]. *Physical Review Letters*, 2009, **102**(18): 181101
- [11] AGUILAR M, AISA D, ALPAT B, *et al.* Precision measurements of the $(e^+ + e^-)$ flux of the primary cosmic rays from 0.5 GeV to 1 TeV with the Alpha magnetic spectrometer on the international space station[J]. *Physical Review Letters*, 2014,

- 113**(22/23/24/25/26/27/28): 221102
- [12] SHEN C S. Pulsars and very high-energy cosmic-ray electrons [J]. *The Astrophysical Journal*, 1970, **162**: L181
 - [13] HOOPER D, BLASI P, SERPICO P D. Pulsars as the sources of high energy cosmic ray positrons[J]. *Journal of Cosmology and Astroparticle Physics*, 2009, **1**: 25
 - [14] YÜKSEL H, KISTLER M D, STANEV T. TeV gamma rays from Geminga and the origin of the GeV positron excess[J]. *Physical Review Letters*, 2009, **103**(5): 051101
 - [15] BERGSTRÖM L, BRINGMANN T, EDSJÖ J. New positron spectral features from supersymmetric dark matter: a way to explain the PAMELA data?[J]. *Physical Review D*, 2008, **78**(10): 103520
 - [16] CIRELLI M, KADASTIK M, RAIDAL M, *et al.* Model-independent implications of the $e^\pm$, $\bar{p}$ cosmic ray spectra on properties of dark matter[J]. *Nuclear Physics B*, 2009, **813**(1/2): 1-21
 - [17] YIN P F, YUAN Q, LIU J, *et al.* PAMELA data and leptonically decaying dark matter[J]. *Physical Review D*, 2009, **79**(2): 023512
 - [18] ABEYSEKARA A U, ALBERT A, ALFARO R, *et al.* Extended gamma-ray sources around pulsars constrain the origin of the positron flux at Earth[J]. *Science*, 2017, **358**(6365): 911-914
 - [19] HOOPER D, GOODENOUGH L. Dark matter annihilation in the Galactic center as seen by the Fermi gamma ray space telescope [J]. *Physics Letters B*, 2011, **697**(5): 412-428
 - [20] CUI M Y, YUAN Q, TSAI Y L S, *et al.* Possible dark matter annihilation signal in the AMS-02 antiproton data[J]. *Physical Review Letters*, 2017, **118**(19): 191101
 - [21] CUOCO A, KRÄMER M, KORSMEIER M. Novel dark matter constraints from antiprotons in light of AMS-02[J]. *Physical Review Letters*, 2017, **118**(19): 191102
 - [22] CHANG J. Dark matter particle explorer: the first Chinese cosmic ray and hard γ -ray detector in Space[J]. *Chinese Journal of Space Science*, 2014, **34**(5): 550-557
 - [23] CHANG J, AMBROSI G, AN Q, *et al.* The dark matter particle explorer mission[J]. *Astroparticle Physics*, 2017, **95**: 6-24
 - [24] YU Y H, SUN Z Y, SU H, *et al.* The plastic scintillator detector for DAMPE[J]. *Astroparticle Physics*, 2017, **94**: 1-10
 - [25] AZZZARELLO P, AMBROSI G, ASFANDIYAROV R, *et al.* The DAMPE silicon-tungsten tracker[J]. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 2016, **831**: 378-384
 - [26] ZHANG Z Y, ZHANG Y L, DONG J N, *et al.* Design of a high dynamic range photomultiplier base board for the BGO ECAL of DAMPE[J]. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 2015, **780**: 21-26
 - [27] HE M, MA T, CHANG J, *et al.* GEANT4 simulation of neutron detector for DAMPE[J]. *Acta Astronomica Sinica*, 2016, **57**(1): 1-8 (何明, 马涛, 常进, 等. 暗物质粒子探测卫星中子探测器的 GEANT4 模拟 [J]. *天文学报*, 2016, **57**(1): 1-8)
 - [28] AMBROSI G, AN Q, ASFANDIYAROV R, *et al.* The on-orbit calibration of DArK matter particle explorer[J]. *Astroparticle Physics*, 2019, **106**: 18-34
 - [29] DONG T K, ZHANG Y P, MA P X, *et al.* Charge measurement of cosmic ray nuclei with the plastic scintillator detector of DAMPE[J]. *Astroparticle Physics*, 2019, **105**: 31-36
 - [30] YUE C, ZANG J J, DONG T K, *et al.* A parameterized energy correction method for electromagnetic showers in BGO-ECAL of DAMPE[J]. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 2017, **856**: 11-16
 - [31] DAMPE Collaboration. Direct detection of a break in the teraelectronvolt cosmic-ray spectrum of electrons and positrons[J]. *Nature*, 2017, **552**(7683): 63-66
 - [32] CHANG J. On the detection and identification of cosmic gamma-rays in a cosmic ray detector[C]//26th International Cosmic Ray Conference. Salt Lake City: ICRC, 1999: 37
 - [33] CHANG J, ADAMS J H, AHN H S, *et al.* Resolving electrons from protons in ATIC[J]. *Advances in Space Research*, 2008, **42**(3): 431-436
 - [34] HUANG Y Y, MA T, YUE C, *et al.* Calibration and performance of the neutron detector onboard of the DAMPE mission[J]. *Research in Astronomy and Astrophysics*, 2020, **20**(9): 153
 - [35] DAMPE Collaboration. Detection of spectral hardenings in cosmic-ray boron-to-carbon and boron-to-oxygen flux ratios with DAMPE[J]. *Science Bulletin*, 2022, **67**(21): 2162-2166
 - [36] HERD Collaboration. The high energy cosmic-radiation detection (HERD) facility onboard China's Space Station[J]. *Proc. SPIE*, 2014, **9144**: 91440X
 - [37] ENGELMANN J J, FERRANDO P, SOUTOUL A, *et al.* Charge composition and energy spectra of cosmic-ray for elements from Be to Ni - Results from HEAO-3-C2[J]. *Astronomy & Astrophysics*, 1990, **233**: 96-111
 - [38] SWORDY S P, MUELLER D, MEYER P, *et al.* Relative abundances of secondary and primary cosmic rays at high energies[J]. *The Astrophysical Journal*, 1990, **349**: 625
 - [39] PANOV A D, SOKOLSKAYA N V, ADAMS J H, *et al.* Relative abundances of cosmic ray nuclei B-C-N-O in the energy region from 10 GeV/n to 300 GeV/n. Results from ATIC-2 (the science flight of ATIC)[C]//30th International Cosmic Ray Conference. Merida: ICRC, 2007: 3
 - [40] AHN H S, ALLISON P S, BAGLIESI M G, *et al.* Measurements of cosmic-ray secondary nuclei at high energies with the first flight of the CREAM balloon-borne experiment[J]. *Astroparticle Physics*, 2008, **30**(3): 133-141
 - [41] OBERMEIER A, AVE M, BOYLE P, *et al.* Energy spectra of primary and secondary cosmic-ray nuclei measured with TRACER[J]. *The Astrophysical Journal*, 2011, **742**(1): 14
 - [42] ADRIANI O, BARBARINO G C, BAZILEVSKAYA G A, *et al.* Measurement of boron and carbon fluxes in cosmic rays with the PAMELA experiment[J]. *The Astrophysical Journal*, 2014, **791**(2): 93
 - [43] GREBENYUK V, KARMANOV D, KOVALEV I, *et al.* Secondary cosmic rays in the NUCLEON space experiment[J]. *Advances in Space Research*, 2019, **64**(12): 2559-2563
 - [44] AGUILAR M, ALI CAVASONZA L A, AMBROSI G, *et al.* The Alpha Magnetic Spectrometer (AMS) on the International

- Space Station: Part II — results from the first seven years[J]. *Physics Reports*, 2021, **894**: 1-116
- [45] DAMPE Collaboration. Measurement of the cosmic p+He energy spectrum from 46 GeV to 316 TeV with the DAMPE space mission[OL]. arXiv preprint arXiv: 2304.00137, 2023
- [46] COLLABORATION D, ASFANDIYAROV R, AZZARELLO P, *et al.* (DAMPE Collaboration). Measurement of the cosmic ray proton spectrum from 40 GeV to 100 TeV with the DAMPE satellite[J]. *Science Advances*, 2019, **5**(9): eaax3793
- [47] ALEMANN F, AN Q, AZZARELLO P, *et al.* Measurement of the cosmic ray helium energy spectrum from 70 GeV to 80 TeV with the DAMPE space mission[J]. *Physical Review Letters*, 2021, **126**(20): 201102
- [48] AHN H S, SEO E S, ADAMS J H, *et al.* The energy spectra of protons and helium measured with the ATIC experiment[J]. *Advances in Space Research*, 2006, **37**(10): 1950-1954
- [49] ATKIN E, BULATOV V, DOROKHOV V, *et al.* First results of the cosmic ray NUCLEON experiment[J]. *Journal of Cosmology and Astroparticle Physics*, 2017, **2017**: 20
- [50] AHN H S, ALLISON P, BAGLIESI M G, *et al.* Discrepant hardening observed in cosmic-ray elemental spectra[J]. *The Astrophysical Journal Letters*, 2010, **714**(1): L89-L93
- [51] BARTOLI B, BERNARDINI P, BI X J, *et al.* Knee of the cosmic hydrogen and helium spectrum below 1 PeV measured by ARGO-YBJ and a Cherenkov telescope of LHAASO[J]. *Physical Review D*, 2015, **92**(9): 092005
- [52] ALBERT A, ALFARO R, ALVAREZ C, *et al.* Cosmic ray spectrum of protons plus helium nuclei between 6 and 158 TeV from HAWC data[J]. *Physical Review D*, 2022, **105**(6): 063021
- [53] ANTONI T, APEL W D, BADEA A F, *et al.* KASCADE measurements of energy spectra for elemental groups of cosmic rays: results and open problems[J]. *Astroparticle Physics*, 2005, **24**(1/2): 1-25
- [54] EAS-TOP Collaboration. The cosmic ray proton, helium and CNO fluxes in the 100 TeV energy region from TeV muons and EAS atmospheric Cherenkov light observations of MACRO and EAS-TOP[J]. *Astroparticle Physics*, 2004, **21**(3): 223-240
- [55] ALEMANN F, ALTOMARE C, AN Q, *et al.* Search for relativistic fractionally charged particles in space[J]. *Physical Review D*, 2022, **106**(6): 063026
- [56] SBARRA C, CASADEI D, BROCCO L, *et al.* Search for fractional charges in cosmic rays with Ams[OL]. arXiv preprint arXiv: astro-ph/0304192, 2003
- [57] FUKU H, TASAKI Y, ABE K, *et al.* Search for fractionally charged particles in cosmic rays with the BESS spectrometer[J]. *Advances in Space Research*, 2008, **41**(12): 2050-2055
- [58] MORI M, OYAMA Y, SUZUKI A, *et al.* Search for fractionally charged particles in Kamiokande II[J]. *Physical Review D*, 1991, **43**(9): 2843-2846
- [59] AGLIETTA M, ANTONIOLI P, BADINO G, *et al.* Search for fractionally charged particles in the Mont Blanc LSD scintillation detector[J]. *Astroparticle Physics*, 1994, **2**(1): 29-34
- [60] AMBROSIO M, ANTOLINI R, AURIEMMA G, *et al.* Search for lightly ionizing particles with the MACRO detector[J]. *Physical Review D*, 2000, **62**(5): 052003
- [61] AGNESE R, ANDERSON A, BALAKISHIYEVA D, *et al.* First direct limits on lightly ionizing particles with electric charge less than $e/6$ [J]. *Physical Review Letters*, 2015, **114**(11): 111302
- [62] ALVIS S I, ARNQUIST I, AVIGNONE F, *et al.* First limit on the direct detection of lightly ionizing particles for electric charge as low as $e/1000$ with the majorana demonstrator[J]. *Physical Review Letters*, 2018, **120**(21): 211804
- [63] DAMPE Collaboration. Search for gamma-ray spectral lines with the DArk Matter Particle Explorer[J]. *Science Bulletin*, 2022, **67**(7): 679-684
- [64] ACKERMANN M, AJELLO M, ALBERT A, *et al.* Updated search for spectral lines from Galactic dark matter interactions with pass 8 data from the Fermi Large Area Telescope[J]. *Physical Review D*, 2015, **91**(12): 122002
- [65] DAMPE Collaboration. Point-like source catalog observed by DAMPE[C]//38th International Cosmic Ray Conference. Nagoya: ICRC, 2023
- [66] LI T P, MA Y Q. Analysis methods for results in gamma-ray astronomy[J]. *The Astrophysical Journal*, 1983, **272**: 317-324
- [67] ABDOLLAHI S, ACERO F, BALDINI L, *et al.* Incremental fermi large area telescope fourth source catalog[J]. *The Astrophysical Journal Supplement Series*, 2022, **260**(2): 53
- [68] Shen Z Q. Analysis of the galactic center excess with DAMPE [C]//38th International Cosmic Ray Conference. Nagoya: ICRC, 2023
- [69] DAYLAN T, FINKBEINER D P, HOOPER D, *et al.* The characterization of the gamma-ray signal from the central Milky Way: a case for annihilating dark matter[J]. *Physics of the Dark Universe*, 2016, **12**: 1-23
- [70] CALORE F, CHOLIS I, WENIGER C. Background model systematics for the Fermi GeV excess[J]. *Journal of Cosmology and Astroparticle Physics*, 2015, **2015**: 038
- [71] CHOLIS I, ZHONG Y M, MCDERMOTT S D, *et al.* Return of the templates: revisiting the Galactic Center excess with multi-messenger observations[J]. *Physical Review D*, 2022, **105**(10): 103023

Author Biography



CHANG Jin Male, born in 1966, chair scientist of the DArk Matter Particle Explorer (DAMPE, also called as “WuKong”) project. He has long been engaged in the detections and researches of gamma ray emission and high energy charged particles that come from outer space.
E-mail: chang@pmo.ac.cn